CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS FOR ANIMAL TREATMENT ONLY

Moxiclear®

For Kittens and Small Cats up to 4 kg / For Cats over 4 kg

100 g/L IMIDACLOPRID 10 g/L MOXIDECTIN

Once a month topical flea and worm treatment

- Stops fleas feeding in 3-5 minutes
- Prevents heartworm and controls intestinal worms and larvae
- Convenient, easy to apply
- Water resistant formulation

For:

- the treatment and prevention of fleas (Ctenocephalides spp) on kittens, cats (small cats up to 4 kg and cats over 4 kg) and ferrets. Larval flea stages are killed in the surroundings of MOXICLEAR® treated pets. Reduces the incidence of Flea Allergy Dermatitis (FAD);
- the prevention of heartworm (*Dirofilaria immitis*) infection in kittens, cats (small cats up to 4 kg and cats over 4 kg) and ferrets.

For the treatment and control of:

- roundworms (Toxocara cati, Toxascaris leonina) kills larval (L4), immature and adult worms:
- hookworms (Ancylostoma tubaeforme, Ancylostoma braziliense (this hookworm is usually restricted to tropical areas in Australia) Uncinaria stenocephala), kills larval (L4), immature and adult worms of all species;
- ear mites (Otodectes cynotis);
- lungworm (Aelurostrongylus abstrusus)

in kittens and cats (small cats up to 4 kg and cats over 4 kg).

One application lasts up to a month.

MOXICLEAR® contains the active constituents, Imidacloprid and Moxidectin. It is effective against sensitive strains of all the above listed parasites. Resistance may develop to any chemicals.

Net Contents:

Moxiclear® For Kittens and Small Cats up to 4 kg [1x0.4 mL] [2x0.4 mL] [3x0.4 mL] [6x0.4 mL] pipette(s)

Moxiclear® For Cats Over 4 kg

[1x0.8 mL] [2x0.8 mL][3x0.8 mL][6x0.8 mL] pipette(s)

It is illegal to sell individual pipette(s) separately

Directions for Use: PRECAUTIONS

Use with caution in sick, debilitated or underweight animals. MOXICLEAR® may be safely applied to heartworm infected cats therefore there is no absolute requirement for a heartworm test before commencing treatment. However a diagnostic test for heartworm prior to commencement of treatment is recommended to ensure the optimum health of cats.

Take care to avoid the product coming into contact with the eyes or mouth of the animal. Ingestion by the recipient or in-contact animals should be prevented.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effect have been observed with imidacloprid or moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating cats. However, the safety of MOXICLEAR® has not been established during pregnancy and lactation.

DOSAGE AND ADMINISTRATION: Consult a veterinarian before use. Treatment for heartworm should occur regularly at monthly intervals.

If worm problems persist, consult a veterinarian.

DO NOT GIVE PRODUCT BY MOUTH. For external application only.

DOSAGE

For kittens and small cats up to 4 kg bodyweight and ferrets	Use 1 PIPETTE of MOXICLEAR® FOR KITTENS AND SMALL CATS UP TO 4 KG
For cats over 4 kg bodyweight	Use 1 PIPETTE of MOXICLEAR® FOR CATS OVER 4 KG

- Remove the pipette from the outer sachet using scissors or fold along diagonal line to expose nick; tear back at nick.
- 2. Hold the pipette upright
- 3. Tap the narrow part of the pipette to ensure the contents remains within the main body of the pipette.
- 4. Twist or snap back the tip.
- 5. Site of Application

Kittens and small cats up to 4 kg	Cats over 4 kg
1 PIPETTE	1 PIPETTE
Apply on the back of the head	Apply on the back of the head

- 6. The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the pipette on the skin and squeeze the pipette firmly twice to apply the solution directly on the skin. Avoid pet's eyes or mouth.
- Record treatment date and repeat monthly or as directed. MOXICLEAR® remains efficacious following shampoo treatment, swimming or exposure to rain. If washing animal during use of this product use a soap free shampoo.
- Occasionally hair loss may occur at the application site. This is usually a single occurrence and hair will grow back uneventfully.

GENERAL INSTRUCTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

FLEA CONTROL AND PREVENTION

MOXICLEAR® has been shown to kill adult and larval fleas within 20 minutes of contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite the animal to receive a lethal dose. The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with MOXICLEAR® rapidly kills fleas and reduces the incidence of this condition. MOXICLEAR® may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the cat. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product.





It is recommended that all cats in the same environment should be protected by MOXICLEAR® for cats and all dogs with MOXICLEAR® for dogs.

HEARTWORM PREVENTION

Treatment with MOXICLEAR® should start one month before the first exposure to mosquitoes. MOXICLEAR® must be applied at one month intervals during the time of the year when mosquitoes (which carry and transmit the heartworm larvae) are present. In many areas of Australia continuous, year-round heartworm prevention is recommended. To establish a routine, it is recommended that the same day or date be used each month to administer MOXICLEAR®. When replacing another heartworm preventive product, the first treatment with MOXICLEAR® must be given within one month of the last dose of the previous product.

WORM CONTROL

MOXICLEAR® treats and controls the major gastrointestinal nematodes of cats but does not control tapeworms. In order to control these worms use specific agent. Cats can become reinfested with worms but regular use of MOXICLEAR® will remove them.

TREATMENT AND CONTROL OF EAR MITES

For the treatment of ear mites (*Otodectes cynotis*), MOXICLEAR® is highly efficacious as a single dose. Monthly application of MOXICLEAR® will control any subsequent ear mite infestation.

To maximise efficacy thorough cleaning of the ear canal is recommended with a product that is registered for that purpose or as directed by a veterinarian.

TREATMENT AND CONTROL OF LUNGWORM

For the treatment of lungworm (*Aelurostrongylus abstrusus*), MOXICLEAR® is highly efficacious as a single dose. Monthly application of MOXICLEAR® will control any subsequent lungworm infections.

KITTENS

Kittens may be treated with MOXICLEAR® FOR KITTENS AND SMALL CATS UP TO 4 KG from 9 weeks of age. If kittens younger than 9 weeks require flea control they may be safely treated with an alternative imidacloprid only ectoparasiticide.

For control of roundworms and hookworms kittens require worming at 6 and 8 weeks of age. When worming kittens prior to commencing MOXICLEAR®, use a specific worming agent. Once treatment with MOXICLEAR® every month is commenced worms will be continuously controlled.

FFRRFTS

Treatment with MOXICLEAR® FOR KITTENS AND SMALL CATS UP TO 4 KG (i.e. 0.4 mL presentation) is recommended. The treatment of ferrets weighing less than 0.8 kg should be based on a risk-benefit assessment by a veterinarian. If ferrets less than 0.8 kg require flea control they may be safely treated with an alternative imidacloprid only ectoparasiticide. Treatment of ferrets may be repeated at two weekly intervals when flea infestation is particularly heavy.

ADDITIONAL INSTRUCTIONS

MOXICLEAR® has no strong chemical smell. The liquid vehicle in MOXICLEAR® may discolour or damage certain surfaces and fabrics e.g polished floorboards, leather. Exclude the animal from contact with such surfaces until the product dries on the animal's coat.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. If product is in eyes, wash it out immediately with water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or a Poisons Information Centre, Phone Australia 13 11 26.

DISPOSAL

Dispose of used pipette(s) and packaging by wrapping in paper and placing in garbage.

STORAGE

Store below 25°C (Air conditioning) and protect from light. Do not freeze

Distributed by: Norbrook Laboratories Australia PTY LTD A.C.N 080 972 596, Unit 7/15-21 Butler Way

Tullamarine, VIC 3043 Free call: 1800 665 866

PRESENTATION

MOXICLEAR® FOR	Contents	APVMA Approval No:
KITTENS AND SMALL CATS UP TO 4 KG	0.4 mL	83601/120018
CATS OVER 4 KG	0.8 mL	83604/120021

WARNING





Acute Toxicity Oral.
Serious eye irritation.
Acute Toxicity Inhalation.
Specific Target Organ
Toxicity - Repeated
Exposure.
Hazardous to aquatic
environment Short
Term/Acute

Harmful if swallowed. Causes eye irritation. Harmful if inhaled. May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life.

Keep out of reach of children. Do not breathe fumes, mists, vapours or spray. Avoid breathing fumes, mists, vapours or spray. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. Avoid release to the environment. Use personal protective equipment as required.

Get medical advice or attention if you feel unwell. Wash with plenty of soap and water. IF SWALLOWED: Rinse mouth. DO NOT induce vomiting. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice. Collect Spillage. In case of fire, use carbon dioxide, dry chemical, foam, water foq.

