

CAUTION
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
READ SAFETY DIRECTIONS

Moxiclear™

For Puppies and Small Dogs up to 4 kg /
 For Dogs 4-10 kg

100 g/L IMIDACLOPRID
25 g/L MOXIDECTIN

Once a month topical flea and worm treatment

- Stops fleas feeding in 3-5 minutes
- Prevents heartworm and controls intestinal worms and larvae
- Convenient, easy to apply
- Water resistant formulation

For:

- the treatment and prevention of fleas *Ctenocephalides spp.*. Reduces the incidence of Flea Allergy Dermatitis, Larval flea stages are killed in the surroundings of MOXICLEAR™ treated pets;
- the prevention of heartworm *Dirofilaria Immitis* infection;

For the treatment and control of:

- roundworms *Toxocara canis*, *Toxascaris leonina* kills larval (L₁), immature and adult worms;
- hookworms *Ancylostoma caninum*, *Ancylostoma braziliense* (This hookworm is usually restricted to tropical areas in Australia) *Ancylostoma ceylanicum*, *Uncinaria stenocephala*, kills larval (L₁), immature and adult worms of all species;
- whipworm *Trichuris vulpis*;
- sarcoptic mange *Sarcoptes scabiei*;
- ear mites *Otodectes cynotis*;
- lice *Trichodectes canis*, *Linognathus setosus* for up to 6 weeks; in dogs and puppies.

One application lasts for a month.

MOXICLEAR™ contains the active constituents, Imidacloprid and Moxidectin. It is effective against sensitive strains of all the above listed parasites. Resistance may develop to any chemicals.

DOSAGE AND ADMINISTRATION: Consult a veterinarian before use. Treatment for heartworm should occur regularly at monthly intervals. For external application only.

DOSAGE

For Small Dogs and Puppies up to 4 kg bodyweight	Use 1 TUBE of MOXICLEAR™ FOR PUPPIES AND SMALL DOGS UP TO 4 KG
For Dogs 4- 10 kg bodyweight	Use 1 TUBE of MOXICLEAR™ FOR DOGS 4- 10 KG

1. Remove required tube(s) from the package
2. Hold tube upright and pull off the cap.
3. Reverse the cap and push cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Site of Application

Dogs under 25 kg
1 TUBE
Apply between the shoulder blades

6. The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the tube on the skin and squeeze the tube firmly **twice to apply the solution directly on the skin.** Avoid pet's eyes or mouth.
7. Record treatment date and repeat monthly or as directed. MOXICLEAR™ remains efficacious following shampoo treatment, swimming or exposure to rain. If washing animal during use of this product use a soap free shampoo.
8. Occasionally hair loss may occur at the application site. This is usually a single occurrence and hair will grow back uneventfully.

GENERAL INSTRUCTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION DO NOT GIVE PRODUCT BY MOUTH

FLEA CONTROL AND PREVENTION

MOXICLEAR™ has been shown to kill adult and larval fleas within 20 minutes of contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite the animal to receive a lethal dose. The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with MOXICLEAR™ rapidly kills fleas and reduces the incidence of this condition. MOXICLEAR™ may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the dog. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product. It is recommended that all dogs in the same environment should be protected by MOXICLEAR™ for dogs and all cats with MOXICLEAR™ for cats.

HEARTWORM PREVENTION

Treatment with MOXICLEAR™ should start one month before the first exposure to mosquitoes. MOXICLEAR™ must be applied at one month intervals during the time of the year when mosquitoes (which carry and transmit the heartworm larvae) are present. In many areas of Australia continuous, year-round heartworm prevention is recommended. To establish a routine, it is recommended that the same day or date be used each month to administer MOXICLEAR™. When replacing another heartworm preventive product, the first treatment with MOXICLEAR™ must be given within one month of the last dose of the previous product.

WORM CONTROL

MOXICLEAR™ treats and controls the major gastrointestinal nematodes of dogs but does not control tapeworms. In order to control tapeworms use appropriate products. Dogs can become reinfested with worms but regular use of MOXICLEAR™ will remove them. **If worm problems persist consult a veterinarian.**

TREATMENT AND CONTROL OF SARCOPTIC MANGE

MOXICLEAR™ has shown to eliminate 100% of sarcoptic mites (*Sarcoptes scabiei*) after a single dose. A second dose of MOXICLEAR™ a month later will result in continued improvement of the clinical signs associated with sarcoptic mange. Monthly application of MOXICLEAR™ will control any subsequent sarcoptic mite infestation.

TREATMENT AND CONTROL OF EAR MITES

For the treatment of ear mites (*Otodectes cynotis*), MOXICLEAR™ is highly efficacious as a single dose. Monthly application of MOXICLEAR™ will control any subsequent ear mite infestation. To maximise efficacy thorough cleaning of the ear canal is recommended with a product that is registered for that purpose or as directed by a veterinarian.

PUPPIES

Puppies may be treated with MOXICLEAR™ from 7 weeks of age. If puppies younger than 7 weeks require flea control they may be safely treated with MOXICLEAR™ FOR PUPPIES AND SMALL DOGS UP TO 4 KG. For control of roundworms and hookworms puppies require worming at **2, 4 and 8 weeks** of age. When worming puppies prior to commencing MOXICLEAR™ use a specific worming agent. Once treatment with MOXICLEAR™ every month is commenced worms will be continuously controlled.

PRECAUTIONS

Do NOT apply to cats. Use with caution in sick, debilitated or underweight animals. MOXICLEAR™ may be safely applied to heartworm infected dogs therefore there is no absolute requirement for a heartworm test before commencing treatment. However a diagnostic test for heartworm prior to commencement of treatment is recommended to ensure the optimum health of dogs.

MOXICLEAR™ has no strong chemical smell. The liquid vehicle in MOXICLEAR™ may discolour or damage certain surfaces and fabrics e.g polished floorboards, leather. Exclude the animal from contact with such surfaces until the product dries on the animal's coat.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effect have been observed with imidacloprid or moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating dogs. However, the safety of MOXICLEAR™ has not been established during pregnancy and lactation.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. If products in eyes, wash it out immediately with water. Wash hands after use.

WARNING



Acute Toxicity Oral.
Serious eye irritation.
Acute Toxicity Inhalation.
Specific Target Organ Toxicity - Repeated Exposure.
Hazardous to aquatic environment Short Term/Acute.

Harmful if swallowed. Causes eye irritation. Harmful if inhaled. May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life.

Keep out of reach of children. Do not breathe fumes, mists, vapours or spray. Avoid breathing fumes, mists, vapours or spray. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. Avoid release to the environment. Use personal protective equipment as required.

Get medical advice or attention if you feel unwell. Wash with plenty of soap and water. IF SWALLOWED: Rinse mouth. DO NOT induce vomiting. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice. Collect Spillage. In case of fire, use carbon dioxide, dry chemical, foam, water fog.

FIRST AID

If poisoning occurs, contact a doctor or a Poisons Information Centre. Phone Australia 131126.

DISPOSAL

Dispose of used tubes and packaging by wrapping in paper and placing in garbage.

PRESENTATION

MOXICLEAR™ FOR	Contents	APVMA Approval No:
Puppies and Small Dogs up to 4 kg	0.4 mL	83619/115813
Dogs 4-10 kg	1 mL	83620/115827

STORAGE

Store below 25°C (Air conditioning) and protect from light. Do not freeze.

Norbroad Laboratories Australia PTY LTD
A.C.N 080 972 596,
Unit 7/15-21 Butler Way
Tullamarine, VIC 3043
Free call: 1800 665 866



Norbroad®